arXiv:1701.00063v2 [hep-th] 23 Jan 2017

January 24, 2017

Stress Tensor Correlators of CCFT$_2$ using Flat-Space Holography

Mohammad Asadi, Omid Baghchesaraei, Reza Fareghbal

Department of Physics, Shahid Beheshti University, G.C., Evin, Tehran 19839, Iran.

m_asadi@sbu.ac.ir ,omidbaghchesaraei@gmail.com, r_fareghbal@sbu.ac.ir

Abstract

We use correspondence between three dimensional asymptotically flat spacetimes and two dimensional contracted conformal field theories (CCFT) to derive the stress tensor correlators of CCFT$_2$. In the gravity side we use the metric formulation instead of the Chern-Simons formulation of three dimensional gravity. This method can also be used for four dimensional case where there is no Chern-Simons formulation for the bulk theory.

1 Introduction

Extending gauge/gravity duality beyond the AdS/CFT correspondence requires that one proposes appropriate dual field theory for the spacetimes which are not asymptotically AdS. One of the potential candidates is asymptotically flat spacetimes. These spacetimes are given by vanishing cosmological constant limit of the asymptotically AdS counterparts. This connection in the gravity side may hint one to propose a dual field theory for the asymptotically flat spacetimes. One of the proposals which links the flat-space limit in the bulk side to the ultra-relativistic limit of the boundary theory, was put forward in [1]-[2]. This proposal which we henceforth call Flat/CCFT, suggests a holographic connection between the asymptotically flat spacetimes in $(d+1)$-dimensions and contracted conformal filed theories (CCFT) in d-dimensions.

A CCFT is given by taking ultra-relativistic limit of the corresponding CFT. In the ultra-relativistic limit the speed of light approches zero and in this singular limit, the symmetries of the theory are not Poincare symmetry. In two dimensions, the contracted conformal algebra is given by Inonu-Wigner contraction of two copies of the Virasoro algebra. Starting with a CFT_2, the contracted algebra is obtained by using generators of the Virasoro algebra and then contracting time-coordinate [2]. The ultra-relativistic limit of the conformal algebra is the opposite of the non-relativistic limit which gives rise to Galilean conformal algebra(GCA)[3]. In two dimensions, these two algebras are isomorphic but in higher dimensions they are different.

A symmetry similar to the contracted conformal symmetry also appears as the asymptotic symmetry of the asymptotically flat spacetimes[4]-[8]. This symmetry which is called BMS symmetry, is infinite dimensional for three and four dimensions. Taking the flat-space limit of the generators of the AdS asymptotic symmetry leads to the generators of the BMS algebra[9]. Thus it is plausible to propose that the ultra- relativistic limit of the CFT is indeed, the dual of the flat-space limit in the asymptotically AdS spacetimes. This idea is used in [1]-[2] where a holographic duality between the asymptotically flat spacetimes and CCFTs is proposed.

Holographic calculation of the stress tensor correlators is a good check for the correctness of the correspondence between a field theory and a gravitational dual theory. It is well-known that the correlation functions of the operators in CFTs have universal forms. One of the success of the AdS/CFT correspondence is its proposed method for the holographic calculation of these correlators.

Similar to the AdS/CFT correspondence, the correlation functions of the operators in a CCFT must have a dual description in the asymptotically flat spacetimes. There are two plausible ways to establish a dictionary which relates calculations in the two sides of the duality. One can ignore the AdS/CFT correspondence and consider Flat/CCFT in its own right or one can take the

appropriate limit of the calculations of the AdS/CFT correspondence. Both of these methods have been invoked and the results have been consistent, so far.

Calculating the stress tensor of CCFT by using flat-space holography was carried out for the first time in [10]. The method used to find the stress tensor of $CCFT_2$, is taking the appropriate limit of the AdS/CFT computations. on the other hand, in [11], a direct method is invoked which yields the correlators of $CCFT_2$. However, the holographic calculations of correlation functions in the gravity side just performed by using the Chern-Simons formulation of three dimensional flat-space gravity. Generalizing such a correspondence to the higher dimensional cases which there is no Chern-Simons formulation for the gravity theory, necessitates the metric formulation of such a calculations.

In the present paper we use the metric formulation of three dimensional gravity in order to calculate the stress tensor correlators via holography. The fact that stress tensor of a field theory can be used to find the conserved charges of the symmetries, is employed to derive an expression for the stress tensor components in terms of the conserved charges. Then the Flat/CCFT proposal is used and the charges are substituted by results in the literature, found directly in the flat spacetimes. Our results in this paper are consistent with [10]. This method has also been used previously for the quasi-local stress tensor of the Kerr black hole [12] and the results are consistent with the ones obtained through taking the flat-space limit.

To calculate the higher-point correlation functions, we make use of invariance of the correlators under the action of the global part of BMS_3 algebra. We track this invariance back to the gravity side and find a general expression for all of the non-zero stress tensor correlator. Our results also confirm the idea that the symmetry algebra of $CCFT_2$ is so rich that it dictates a universal form for the correlators. Another non-trivial point in our calculation is assuming a non-symmetric stress tensor for the CCFTs. Our investigations in the gravity side show that a covariant conservation formula requires a non-symmetric stress tensor. The fact that CCFTs do not exhibit Poincare symmetry helps us avoid any inconsistencies. Our calculations in the present paper provide yet another confirmation for the fact that asymptotically flat spacetimes do have holographic duals which are CCFTs living in one less dimension.

In section two we introduce the stress tensor of $CCFT_2$ by using holographic method and metric formulation of three dimensional gravity. In section three we calculate the p-point functions of the stress tensor by using holography. The last section, section 4, is devoted to discussion and directions for possible future investigations.

2 Stress Tensor of CCFT

Our goal is to calculate the correlation functions of the $CCFT_2$ stress tensor. In the first step we need to introduce the stress tensor. According to our convention, a $CCFT_2$ is a theory which is defined by the following infinite dimensional symmetry:

$$\begin{aligned}[L_m, L_n] &= (m-n)L_{n+m} + C_L m(m^2-1)\delta_{m+n,0}, \\ [L_m, M_n] &= (m-n)M_{n+m} + C_M m(m^2-1)\delta_{m+n,0},\end{aligned} \tag{2.1}$$

where n and m can take any integer values. Similar to CFT_2, one may expect that the above infinite dimensional symmetry yields some universal results which are independent of the underlying action. The Algebra (2.1) is given by Inonu-Wigner contraction of the Virasoro algebra. Thus, one may consider $CCFT_2$ as a contracted theory obtained from a parent CFT. There are two possible contractions of the Virasoro algebra which lead to (2.1), a Non-relativistic and an ultra relativistic contraction. The first one which is given by taking very large limit of the light speed, corresponds to scaling $x \to \epsilon x$ and $\epsilon \to 0$. On the other hand the ultra relativistic contraction is obtained by vanishing limit of light speed or equivalently scaling $t \to \epsilon t$ and $\epsilon \to 0$. In two dimensions both of the non-relativistic and ultra relativistic contractions of the Virasoro algebra give rise to the same algebra as in (2.1). However in general , by CCFT we mean a theory which its symmetry is given by the ultra-relativistic limit. The non-relativistic limit yields Galilean conformal algebra (GCA) which is interesting in its own[3].

We suppose that $CCFT_2$ lives on a cylinder with metric

$$ds^2 = -du^2 + R^2 d\phi^2 \tag{2.2}$$

where R is the radius of the cylinder which will be fixed later when we use the holographic dictionary. Our starting point for finding the stress tensor of CCFT is the formula which gives the conserved charges of symmetry generators ξ. Using (2.2) we can write

$$Q_\xi = R\int_0^{2\pi} d\phi J^u = R\int_0^{2\pi} d\phi T^{u\mu}\xi_\mu, \tag{2.3}$$

where J^μ is the symmetry current and $T^{\mu\nu}$ is the stress tensor. Here, we do not impose any conditions on the components of the stress tensor. For a CCFT that lives on the cylinder one can introduce a representation for the generators of (2.1) as below

$$L_n = ie^{in\phi}\left(\partial_\phi + inu\partial_u\right), \qquad M_n = ie^{in\phi}\partial_u. \tag{2.4}$$

Thus we can write

$$Q_{M_n} = -iR\int_0^{2\pi} d\phi\, e^{in\phi}\, T^{uu},$$
$$Q_{L_n} = R\int_0^{2\pi} d\phi\, e^{in\phi}\, \left(nuT^{uu} + iR^2T^{u\phi}\right). \tag{2.5}$$

Using the orthogonality condition of Fourier modes, we can find T^{uu} and $T^{u\phi}$ from (2.5) as

$$T^{uu} = \frac{i}{2\pi R}\sum_n Q_{M_n} e^{-in\phi}$$
$$T^{u\phi} = \frac{-i}{2\pi R^3}\sum_n e^{-in\phi}\left(Q_{L_n} - iunQ_{M_n}\right) \tag{2.6}$$

The other components must be determined by using the conservation and traceless-ness conditions. However, in order to check the above calculations and find other components we make use of Flat/CCFT proposal and first do a holographic calculation.

2.1 Holographic calculation using Flat/CCFT correspondence

The calculations in the previous section are pure field theoretic ones and we merely defined a two dimensional field theory by its symmetries. However, as is proposed in [1]-[2] this two dimensional field theory has a holographic dual theory. The dual theory is three dimensional gravity in asymptotically flat backgrounds. The asymptotic symmetries of such a spacetimes at null infinity is known as BMS_3 symmetry which is isomorphic to (2.1). Thus we can find an interpretation for the charges Q_{M_n} and Q_{L_n} in the bulk side as the charges corresponding to the asymptotic symmetry generators. To be precise, let us consider a set of asymptotically flat spacetimes which transforms back into itself under the action of asymptotic symmetry generators. In a particular coordinate systems, known as BMS coordinates, the generic form of the asymptotically flat spacestimes with BMS_3 asymptotic symmetry is given by [8]

$$ds^2 = Mdu^2 - 2dudr + 2Ndud\phi + r^2d\phi^2, \tag{2.7}$$

where

$$M = \theta(\phi), \qquad N = \chi(\phi) + \frac{u}{2}\theta'(\phi), \tag{2.8}$$

and $\theta(\phi)$ and $\chi(\phi)$ are arbitrary functions of ϕ coordinate. u is known as *retarded time* where for the Minkowski spacetime $u = t - r$. The generators of infinitesimal coordinate transformation, ξ^μ, which preserve form of the metric (2.7) are given by

$$\xi^u = F, \qquad \xi^\phi = Y - \frac{1}{r}\partial_\phi F, \qquad \xi^r = -r\partial_\phi Y + \partial_\phi^2 F - \frac{1}{r}N\partial_\phi F, \tag{2.9}$$

where

$$Y = Y(\phi), \qquad F = T(\phi) + uY'(\phi), \tag{2.10}$$

$Y(\phi)$ and $T(\phi)$ are arbitrary functions. L_n and M_n which are defined by

$$L_n = \xi(Y = ie^{in\phi}, T = 0), \qquad M_n = \xi(Y = 0, T = ie^{in\phi}), \tag{2.11}$$

satisfy the algebra (2.1) at large r. The corresponding charges of L_n and M_n can be computed by various methods. They are given by *covariant phase space* method [13],[8] as[1]

$$\begin{aligned} Q_{M_n} &= \frac{i}{16\pi G}\int_0^{2\pi} d\phi\, e^{in\phi}\theta(\phi) + \frac{i}{8G}\delta^0_n, \\ Q_{L_n} &= \frac{i}{8\pi G}\int_0^{2\pi} d\phi\, e^{in\phi}\chi(\phi). \end{aligned} \tag{2.12}$$

The shift in the first line of (2.12) is necessary in order to the Poisson bracket of the charges produce the correct coefficient for the central term in the algebra (2.1). The interesting point here is that with this shift of charges we have $Q_{M_0} = Q_{L_0} = 0$ for the Minkowski metric.

Substituting (2.12) in (2.6) one can find components of the stress tensor as follows:

$$\begin{aligned} T^{uu} &= -\frac{1}{16\pi GR}\left(1 + \theta(\phi)\right), \\ T^{u\phi} &= \frac{1}{8\pi GR^3}\left(\chi(\phi) + \frac{u}{2}\theta'(\phi)\right). \end{aligned} \tag{2.13}$$

This result is consistent with those of [10] where components of the stress tensor are calculated through taking flat-space limit from the quasi-local stress tensor of the asymptotically AdS space-times. Moreover, we find the same results as in [11] if M and N in [11] are identified as T_{uu} and $T_{u\phi}$ components of the stress tensor. We have not fixed the constant R in the above calculations,yet. It can be done through relating the constant term in the uu component of the stress tensor with the central charges of (2.1).

By assuming a standard conservation formula for the components of the stress tensor one arrives at

$$\partial_u T^{u\phi} + \partial_\phi T^{\phi\phi} = 0. \tag{2.14}$$

Thus using (2.13) we can determine $T^{\phi\phi}$ as

$$T^{\phi\phi} = -\frac{\theta(\phi)}{16\pi GR^3} + K, \tag{2.15}$$

where K is a constant of integration. If we also impose a traceless-ness condition $T^\mu_\mu = 0$ for the stress tensor, K is determined and we have

$$T^{\phi\phi} = -\frac{1}{16\pi GR^3}\left(1 + \theta(\phi)\right). \tag{2.16}$$

[1] The calculation of surface charges in [8] has been done at the circle at infinity. Moreover it is assumed that the background line element which is used to raise and lower indices is Minkowski, $ds^2 = -du^2 - 2dudr + r^2 d\phi^2$.

From (2.13) it is clear that the conservation equation,

$$\partial_u T^{uu} + \partial_\phi T^{\phi u} = 0, \tag{2.17}$$

is not satisfied for a symmetric stress tensor, i.e. $T^{u\phi} = T^{\phi u}$. One possible way to overcome this obstacle is assuming a new conservation equation as $\partial_u T^{uu} = 0$ [10]. However, if we want to write the conservation formula in a covariant way, there is a possibility of assuming non-symmetric stress tensors for the CCFTs. If we implement a non-symmetric stress tensor (similar to the case in [14]) such that $T^{u\phi}$ is non-zero and is given by (2.13) but $T^{\phi u} = 0$ then the holographic calculations results in the standard conservation equation, $\nabla_\mu T^{\mu\nu} = 0$ for the CCFT. The fact that CCFTs are not Poincare invariant theories makes this assumption reliable. We should note again that all of these results are consequences of accepting a holographic duality between CCFTs and asymptotically flat spacetimes. In summary, we have

$$\begin{aligned}
T_{uu} &= -\frac{1}{16\pi GR}\left(1+\theta(\phi)\right), \\
T_{u\phi} &= -\frac{1}{8\pi GR}\left(\chi(\phi) + \frac{u}{2}\theta'(\phi)\right), \\
T_{\phi\phi} &= -\frac{R}{16\pi G}\left(1+\theta(\phi)\right), \\
T_{\phi u} &= 0.
\end{aligned} \tag{2.18}$$

3 Correlators of stress Tensor

In this section we use the results of the previous sections to calculate the correlation functions of CCFT_2. To do so, we assume that these functions are invariant under the global part of the two dimensional symmetry algebra. For the two dimensional theory, whose symmetry is given by (2.1), the global part is generated by $\{L_0, L_{\pm 1}, M_0, M_{\pm 1}\}$. According to (2.18), the holographic calculations yield the components of stress tensor in terms of two functions $\theta(\phi)$ and $\chi(\phi)$. When we fix these functions in the gravity side, the asymptotically flat solution is completely determined. An infinitesimal coordinate transformation generated by (2.9) changes these functions to $\theta+\delta\theta$ and $\chi+\delta\chi$. The infinitesimal changes of the functions can be calculated by using the Lie derivative of the metric components and expressing them in such a way that the generic form (2.7) is preserved. We arrive at[]

$$\begin{aligned}
\delta_\xi \theta &= Y\theta' + 2Y'\theta - 2Y''', \\
\delta_\xi \chi &= \frac{1}{2}T\theta' + Y\chi' + 2Y'\chi + T'\theta - T'''.
\end{aligned} \tag{3.1}$$

We apply (3.1) in the gravity side to find the variation of the stress tensor in the boundary. Using (2.18) and (3.1) and imposing conditions

$$\delta_{M_n}\langle T_{ij}\rangle = 0, \qquad \delta_{L_n}\langle T_{ij}\rangle = 0, \qquad n = 0, \pm 1 \tag{3.2}$$

result in

$$\langle T_{ij}\rangle = 0, \tag{3.3}$$

as expected.

We can also use (2.18) and (3.1) to calculate higher-point functions. Since according to (2.18), $T_{\phi\phi}$ is the same as T_{uu} upto a overall factor, its correlation functions with the other components are similar to the correlation functions of T_{uu}. Similar to the one-point functions, we want to determine the p-point functions by imposing

$$\delta_{M_n}\langle T^1_{ij}\cdots T^p_{kl}\rangle = 0, \qquad \delta_{L_n}\langle T^1_{ij}\cdots T^p_{kl}\rangle = 0, \qquad n = 0, \pm 1 \tag{3.4}$$

where $T^l_{ij} = T_{ij}(u_l, \phi_l)$.

If we define $\beta(\phi) = \theta(\phi) + 1$ then the uu and $\phi\phi$ components of the stress tensor will be proportional to $\beta(\phi)$. For $n = 0, \pm 1$, equations (2.11) and (3.1) yield the following variations:

$$\begin{aligned} \delta_{M_n}\beta &= 0, \qquad & \delta_{L_n}\beta &= e^{in\phi}\left(i\partial_\phi\beta - 2n\beta\right), \\ \delta_{M_n}\chi &= \frac{1}{2}e^{in\phi}\left(i\partial_\phi\beta - 2n\beta\right), \qquad & \delta_{L_n}\chi &= e^{in\phi}\left(i\partial_\phi\chi - 2n\chi\right). \end{aligned} \tag{3.5}$$

It is clear from (3.5) that imposing $\delta_{L_n}\langle T^1_{ij}\cdots T^p_{kl}\rangle = 0$ for $n = 0, \pm 1$ results in equations

$$\sum_{k=1}^{P}\langle X_1\cdots e^{in\phi_k}(i\partial_k - 2n)X_k\cdots X_p\rangle = 0, \tag{3.6}$$

where X_i can be either of $\beta_i = \beta(\phi_i)$ or $\chi_i = \chi(\phi_i)$ and ∂_k indicates derivative with respect to ϕ at point ϕ_k. Thus we conclude that for a given p, all of the p-point functions of β and χ with any numbers of β and χ and any insertion of them have the same functionality of $\{\phi_1, \phi_2, \cdots, \phi_p\}$ but with different overall constant factors. These constants can also be zero which would render some correlation functions to vanish.

The solution to equation (3.6) is given by

$$\langle X_1\cdots X_p\rangle = C\frac{e^{2i\sum_{k=1}^{p}\phi_k}}{\prod_{1\le l<m\le p}\left(e^{i\phi_l} - e^{i\phi_m}\right)^{\frac{4}{p-1}}}, \tag{3.7}$$

where C is a constant which can be zero. We determine C by imposing $\delta_{M_n}\langle T^1_{ij}\cdots T^p_{kl}\rangle = 0$ for $n = 0, \pm 1$. $\langle\chi_1\cdots\chi_p\rangle$ does not appear in any other equation thus we conclude that it is given by the generic equation (3.7). Moreover, from $\delta_{M_n}\langle T^1_{u\phi}\cdots T^p_{u\phi}\rangle = 0$ we find that

$$\langle\beta_1\chi_2\chi_3\cdots\chi_p\rangle = \langle\chi_1\beta_2\chi_3\cdots\chi_p\rangle = \cdots = \langle\chi_1\chi_2\chi_3\cdots\beta_p\rangle. \tag{3.8}$$

However, applying δ_{M_n} to other correlation functions which have at least one T_{uu} reveals that the correlations of β and χ containing more than one β must be zero in order to be consistent with the generic result (3.7). Thus the only non-zero correlation functions of β and χ are

$$\begin{aligned}\langle \chi_1 \cdots \chi_p \rangle &= C_1 \frac{e^{2i\sum_{k=1}^p \phi_k}}{\prod_{1\le l<m\le p}(e^{i\phi_l}-e^{i\phi_m})^{\frac{4}{p-1}}},\\ \langle \beta_1\chi_2\chi_3\cdots\chi_p\rangle = \cdots = \langle \chi_1\chi_2\chi_3\cdots\beta_p\rangle &= C_2 \frac{e^{2i\sum_{k=1}^p \phi_k}}{\prod_{1\le l<m\le p}(e^{i\phi_l}-e^{i\phi_m})^{\frac{4}{p-1}}}\end{aligned} \tag{3.9}$$

where C_1 and C_2 are constants. They must be related to the central charges c_M and c_L of (2.1) but in the gravity side they are only constants of integration. Now using (2.18) and (3.9) we can calculate all of the correlation functions of the stress tensor:

$$\begin{aligned}\langle T^1_{uu} T^2_{u\phi}\cdots T^p_{u\phi}\rangle &\propto C_2 \frac{e^{2i\sum_{k=1}^p \phi_k}}{\prod_{1\le l<m\le p}(e^{i\phi_l}-e^{i\phi_m})^{\frac{4}{p-1}}}\\ \langle T^1_{u\phi} T^2_{u\phi}\cdots T^p_{u\phi}\rangle &\propto \left(C_1 + \frac{C_2}{2}\sum_{k=1}^p u_k\partial_k\right)\frac{e^{2i\sum_{k=1}^p \phi_k}}{\prod_{1\le l<m\le p}(e^{i\phi_l}-e^{i\phi_m})^{\frac{4}{p-1}}}\end{aligned} \tag{3.10}$$

It is clear from (3.10) that all of the p-point functions of $T_{u\phi}$s are given by correlation of one T_{uu} and $p-1$ of $T_{u\phi}$s.

For the two-point functions we find

$$\begin{aligned}\langle T_{uu}(\phi_1)T_{uu}(\phi_2)\rangle &= 0,\\ \langle T_{uu}(\phi_1)T_{u\phi}(u_2,\phi_2)\rangle &\propto C_2\frac{e^{2i(\phi_1+\phi_2)}}{(e^{i\phi_1}-e^{i\phi_2})^4},\\ \langle T_{u\phi}(u_1,\phi_1)T_{u\phi}(u_2,\phi_2)\rangle &\propto C_1\frac{e^{2i(\phi_1+\phi_2)}}{(e^{i\phi_1}-e^{i\phi_2})^4}\left(1+2i\frac{C_2}{C_1}(u_2-u_1)\frac{e^{i\phi_1}+e^{i\phi_2}}{e^{i\phi_1}-e^{i\phi_2}}\right).\end{aligned} \tag{3.11}$$

which is consistent with the results in [11].

4 Discussion

In this paper we calculate all correlation functions of CCFT$_2$ stress tensor by using Flat/CCFT proposal. In the gravity side we make use of the metric formulation of three dimensional gravity. Our method is applicable for higher dimensional cases, as well. The interesting point is that the asymptotic symmetry of four dimensional asymptotically flat spacetimes is also infinite dimensional. The symmetry algebra is known as BMS$_4$ algebra. Using Flat/CCFT correspondence we conclude that CCFT$_3$ also has an infinite dimensional symmetry and therefore one expects universal behaviour for the correlation functions of the operators[15]. Since the conformal symmetry in three dimensions is finite dimensional, it is not clear how to find the correlators by taking

the flat space limit. However, it is possible to generalize our method in this paper to the four dimensional case. If this is done successfully some related interesting questions arise. Among them the issue of entanglemet entropy of $CCFT_3$ which is expected to use the universality of the correlation functions[16]-[18].

Another interesting question in the context of $Flat_3/CCFT_2$ proposal is calculation of higher correlation functions by taking the appropriate limit of the AdS/CFT calculations. This might be done along the lines which was first introduced in [19]. The importance of this problem is that its solution is a necessary step in the path to find holographic renormalization method for the Flat/CCFT correspondence.

Acknowledgements

The authors would like to thank A. Bagchi for useful comments. R. F would like to thank school of particles and accelerator of institute for research in fundamental sciences (IPM) for the research facilities.

References

[1] A. Bagchi, "Correspondence between Asymptotically Flat Spacetimes and Nonrelativistic Conformal Field Theories," Phys. Rev. Lett. **105**, 171601 (2010).

A. Bagchi, "The BMS/GCA correspondence," arXiv:1006.3354 [hep-th].

[2] A. Bagchi and R. Fareghbal, "BMS/GCA Redux: Towards Flatspace Holography from Non-Relativistic Symmetries," JHEP **1210**, 092 (2012) [arXiv:1203.5795 [hep-th]].

[3] A. Bagchi and R. Gopakumar, "Galilean Conformal Algebras and AdS/CFT," JHEP **0907**, 037 (2009) [arXiv:0902.1385 [hep-th]].

[4] H. Bondi, M. G. van der Burg, and A. W. Metzner, "Gravitational waves in general relativity. 7. Waves from axisymmetric isolated systems," *Proc. Roy. Soc. Lond. A* **269** (1962) 21.

R. K. Sachs, "Gravitational waves in general relativity. 8. Waves in asymptotically flat space-times," *Proc. Roy. Soc. Lond. A* **270** (1962) 103.

R. K. Sachs, "Asymptotic symmetries in gravitational theory," *Phys. Rev.* **128** (1962) 2851.

[5] A. Ashtekar, J. Bicak and B. G. Schmidt, "Asymptotic structure of symmetry reduced general relativity," Phys. Rev. D **55**, 669 (1997) [gr-qc/9608042].

[6] G. Barnich and G. Compere, "Classical central extension for asymptotic symmetries at null infinity in three spacetime dimensions," Class. Quant. Grav. **24**, F15 (2007) [gr-qc/0610130].

[7] G. Barnich and C. Troessaert, "Symmetries of asymptotically flat 4 dimensional spacetimes at null infinity revisited," arXiv:0909.2617 [gr-qc].

[8] G. Barnich and C. Troessaert, "Aspects of the BMS/CFT correspondence," JHEP **1005**, 062 (2010) [arXiv:1001.1541 [hep-th]].

[9] G. Barnich, A. Gomberoff and H. A. Gonzalez, "The Flat limit of three dimensional asymptotically anti-de Sitter spacetimes," Phys. Rev. D **86**, 024020 (2012) doi:10.1103/PhysRevD.86.024020 [arXiv:1204.3288 [gr-qc]].

[10] R. Fareghbal and A. Naseh, "Flat-Space Energy-Momentum Tensor from BMS/GCA Correspondence," JHEP **1403**, 005 (2014) [arXiv:1312.2109 [hep-th]].

[11] A. Bagchi, D. Grumiller and W. Merbis, "Stress tensor correlators in three-dimensional gravity," Phys. Rev. D **93**, no. 6, 061502 (2016) doi:10.1103/PhysRevD.93.061502 [arXiv:1507.05620 [hep-th]].

[12] O. Baghchesaraei, R. Fareghbal and Y. Izadi, "Flat-Space Holography and Stress Tensor of Kerr Black Hole," Phys. Lett. B **760**, 713 (2016) doi:10.1016/j.physletb.2016.07.062 [arXiv:1603.04137 [hep-th]].

[13] G. Barnich and F. Brandt, "Covariant theory of asymptotic symmetries, conservation laws and central charges," Nucl. Phys. B **633**, 3 (2002) doi:10.1016/S0550-3213(02)00251-1 [hep-th/0111246].

[14] J. Hartong, "Holographic Reconstruction of 3D Flat Space-Time," JHEP **1610**, 104 (2016) doi:10.1007/JHEP10(2016)104 [arXiv:1511.01387 [hep-th]].

[15] A. Bagchi, R. Basu, A. Kakkar and A. Mehra, "Flat Holography: Aspects of the dual field theory," arXiv:1609.06203 [hep-th].

[16] A. Bagchi, R. Basu, D. Grumiller and M. Riegler, "Entanglement entropy in Galilean conformal field theories and flat holography," Phys. Rev. Lett. **114**, no. 11, 111602 (2015) doi:10.1103/PhysRevLett.114.111602 [arXiv:1410.4089 [hep-th]].

[17] S. M. Hosseini and . Vliz-Osorio, "Gravitational anomalies, entanglement entropy, and flat-space holography," Phys. Rev. D **93**, no. 4, 046005 (2016) doi:10.1103/PhysRevD.93.046005 [arXiv:1507.06625 [hep-th]].

[18] R. Basu and M. Riegler, "Wilson Lines and Holographic Entanglement Entropy in Galilean Conformal Field Theories," Phys. Rev. D **93**, no. 4, 045003 (2016) doi:10.1103/PhysRevD.93.045003 [arXiv:1511.08662 [hep-th]].

[19] R. N. Caldeira Costa, "Aspects of the zero Λ limit in the AdS/CFT correspondence," Phys. Rev. D **90**, no. 10, 104018 (2014) doi:10.1103/PhysRevD.90.104018 [arXiv:1311.7339 [hep-th]].